Validation of the 2014 ESC
Guidelines Risk Prediction
Model for the Primary
Prevention of Sudden Cardiac
Death in Hypertrophic
Cardiomyopathy

Journal club, May 20th, 2015

HCMにおけるICD植え込みの適応 (JCS2012)

Class I

過去に持続性心室頻拍，心室細動，心停止の既往を有する場合

Class IIa

非持続性心室頻拍
突然死の家族歴
失神
左室壁厚30mm以上
運動時の血圧反応異常

のいずれかを認める場合に考慮する

HCMにおけるICD植え込みの適応 (ACC/AHA2011)

Class I
心停止、VF、持続性VTに対する二次予防目的
(Level B)

Class IIa
1) 1親等以内の突然死家族歴
壁厚 ≥ 30 mm
最近の失神 のいずれかを認める場合

2) NSVT or 血圧反応異常
かつ他のリスクあり
(LVOT obstruction、LGE on CMR、
LV apical aneurysm、Malignant mutation(?))
(Level C)

Class IIb
NSVTのみ or 血圧反応異常のみ
(Level C)

HCMにおけるICD植え込みの適応 (ESC2014)

SECONDARY PREVENTION

- Cardiac arrest due to VT or VF
- spontaneous sustained VT causing syncope or haemodynamic compromise

Life expectancy > 1 year

ICD recommended

Class I (Level B)

PRIMARY PREVENTION

Recommended assessment:

History
2D/Doppler echocardiogram
48-hour ambulatory ECG

HCM Risk-SCD variables:

- Age
- Family history of sudden cardiac death
- Unexplained syncope
- Left ventricular outflow gradient^a
- Maximum left ventricular wall thickness^a
- Left atrial diameter^a
- NSVT

HCM Risk-SCD Score

Class IIb or III (Level B)

Class IIb (Level B)

Class IIa (Level B)

The risk of SCD in 5 years

$$\hat{P}_{\text{SCD at 5 years}} = 1 - 0.998^{\exp(\text{Prognostic Index})},$$

where Prognostic Index = $0.15939858 * \text{Maximal wall thickness (mm)} - 0.00294271 * \text{Maximal wall thickness}^2 \text{ (mm}^2\text{)} + 0.0259082 * \text{Left atrial diameter (mm)} + 0.00446131 * \text{Maximal left ventricular outflow tract gradient (mmHg)} + 0.4583082 * \text{Family history SCD} + 0.82639195 * \text{NSVT} + 0.71650361 * \text{Unexplained syncope} - 0.01799934 * \text{Age at clinical evaluation (years)}$.

NRI: net reclassification improvement

Table 3: Reclassification of predicted risk among cases (patients with SCD-event) and controls

	Predicted risk classified downward in new model	Predicted risk not changed in new model	Predicted risk classified upward in new model	Total
<i>2014 vs. 2003 guidelines model</i>				
Cases (SCD-patients), n (%)	0 (0) †	11 (55)	9 (45)*	20
Controls, n (%)	12 (2)*	540 (79)	134 (20) †	686
<i>2014 vs. 2011 guidelines model</i>				
Cases (SCD-patients), n (%)	2 (10) †	12 (60)	6 (30)*	20
Controls, n (%)	75 (11)*	513 (75)	98 (14) †	686

NRI

0.27

(95%CI -0.02-0.57,
p=0.07)

0.16

(95%CI -0.17-0.45,
p=0.2)

* indicates correct reclassifications in the new model. † indicates incorrect reclassifications in the new model. SCD = sudden cardiac death

NNT: number needed to treat

1人のSCDを予防するのに何人ICDを植え込めばよいか

Table 4: ICD implantations and 5-year risk of SCD based on the HCM Risk-SCD model, and 2003 and 2011 risk prediction models

	Patients with SCD		Patients without SCD	
	<i>n</i> = 706 ICD	No ICD	ICD	No ICD
2003 guidelines (≥ 2 established risk factors)	5 (25)	15 (75)	111 (16)	575 (84)
2011 guidelines (≥ 1 established risk factor)	10 (50)	10 (50)	210 (31)	476 (69)
HCM Risk-SCD score $\geq 4\%$	14 (70)	6 (30)	229 (33)	457 (67)

Data are represented as number (percentage). HCM: hypertrophic cardiomyopathy; ICD: implantable cardioverter defibrillator; SCD: sudden cardiac death.

NNT

22

20

17